


# Surface issues for SRNWP

J.-F. Mahfouf, S. Tijm,  
A. Trojakova, M. Best, B. Ritter

Thanks to J.-M. Bettems, D. Mironov, P. Pellerin, J.-L. Roujean, D. Carrer, Y. Seity,  
J.-C. Calvet, J. Cedilnik



# Outline

- Specificities of surface aspects for SRNWP
  - Coupling strategies
  - Modelling aspects
  - Physiographic data bases
  - Data assimilation aspects
  - Validation studies
- 



# Purpose of surface schemes

- Realistic description of momentum, energy and mass (water, CO<sub>2</sub>, aerosols, ...) exchanges between the Earth's surface and the atmospheric turbulent boundary layer.
- Framework of SRNWP :
  - Temporal scales : < ~ few days
  - Spatial scales : few kms



# Main surface types

Bare soil




Vegetation



Snow



Oceans



Ice caps



Sea ice



# Main surface types

Bare soil




Vegetation



Snow



Oceans




Ice caps



Sea ice



Towns



Lakes





# How to characterize surfaces ?

- Fast evolving components ( $< 1$  day) : **modelling** (evolution of prognostic variables)
- Slow evolving components ( $\sim 1$  week) : **data assimilation** and surface analyses
- Static components ( $> 1$  month) : specification though **physiographic data bases**
- Land surface temperature, interception reservoir
- Root zone soil moisture – snow cover - SST
- Soil texture, land cover





# Motivations for collaborations


- Common interest on some aspects –  
Share common experience
- Complexity of surface processes =>  
Benefit from complementary expertise
- Take advantage of European initiatives  
(e.g. EUMETSAT SAF – EUMETNET  
OPERA)
- Share national validation data bases








# Coupling strategies

- Surface externalised from the atmospheric model : to be used by various models (NWP, mesoscale research, climate, hydrology), offline validations and data assimilation
  - Existing platforms : SURFEX and JULES
  - How is consistency achieved between modules (e.g. subgrid heterogeneities, orography, continuity between SBL and ABL ?)
  - How can modularity be achieved (external vs internal coupler) ?
- 




# Internal coupling strategy (SURFEX)



ALMA coupling norm (Polcher et al., 1998; Best et al., 2004)


# Mosaic approach in COSMO



From J.-M. Bettems (Meteo Swiss)

# External coupling strategy

Canadian environmental modelling system





# Current modelling developments


- Description of missing processes in SURFEX (ALADIN/LACE) already available in HIRLAM : snow/forest interactions (workshop in Toulouse), prognostic sea-ice
- New (common) area of development : use of FLake
- Common interest in stable surface boundary layers : CANOPY in SURFEX, QNSE theory in HIRLAM
- What else is relevant for SRNWP (available in SURFEX) : towns, ocean, interactive vegetation, aerosol transport, ... ?



# Lake Fraction in COSMO LM1

Lake-fraction external-parameter field for LM1 domain.

mean: 0.17 std: 0.20 min: 0.02 max: 1.00




0.00 <= unknown 2008010100 0000 0 1 1 DWD ,u/work1/dminonov/vs61/GRIB/FR\_LAKE\_LM1 <= 1.00 correlation(field,filter): 1.000

Lake-fraction external-parameter field for the LM1 numerical domain (DWD) of the NWP model COSMO based on the GLCC data set (<http://edcsns17.cr.usgs.gov/glcc/>) with 30 arc sec resolution, that is ca. 1 km at the equator.

From D. Minonov (DWD)

# FLake in COSMO: Results from Parallel Experiment 5632

## 1 January – 31 December 2006




Lake Balaton, Hungary (mean depth = 3.3 m)

- Black – lake surface temperature from the COSMO-LM SST analysis
- Green – lake surface temperature computed with FLake

From D. Minonov (DWD)


# Impact of urban tiles

Aladin 2006041300+1200 T2m (C) & V10m




**ALADIN**  
 $\Delta x = 9.5$  km

Aro 2006041300+1200 T2m (C) & V10m



**AROME-TEB**  
 $\Delta x = 2.5$  km

Obs 2006041300+1200 Obs T2m (C)






**SYNOP**  
observations

Montpellier


12-h forecast of screen-level temperature



Sea-ice  
thickness



Sensible heat flux ( $\text{W/m}^2$ )




Temperature ( $^\circ\text{C}$ )

From P. Pellerin (MSC)



# Physiographic databases


- Resolution issues : soil texture global products (FAO) at 10 km
  - Availability of high resolution land cover maps over Europe : CORINE2000 (100 m)
  - Interest for regional data sets vs global data sets (LAM outside Europe) ?
  - Land cover maps = climatologies => use of real-time satellite products
  - Databases for new surface types : towns, lakes ?
- 







# ECOCLIMAP 2

- Revised land cover climatology at Météo-France
  - Higher resolution and improved quality of land cover maps
  - Multi-year data availability
  - Improved method for ecosystem classification
- 

CORINE2000 – 100m





GLC2000 – 1 km




1-km merged  
land cover map

From J.L. Roujean (MF)




Crop cover



NDVI SPOT/VGT

Automatic  
classification



Crop  
ecosystems


From  
J.-L. Roujean (MF)





# ECOCLIMAP II

Resulting land cover map: 305 ecosystems



Look-up tables  
for internal  
model  
parameters



# Data assimilation aspects (1)

- Better usage of remote-sensing data (available over Europe)
  - SST and Sea-ice : OSI-SAF – MERSEA - OSTIA
  - Snow analysis : snow cover / snow water equivalent (LandSAF, MODIS)
  - Soil moisture analysis : satellite derived products (ASCAT, AMSR-E, SMOS)
  - Vegetation properties : albedo – LAI (LandSAF, MODIS)
  - Radiative forcing : downward fluxes (LandSAF)
  - Precipitation forcing : radar networks (OPERA)
- 







## Data assimilation aspects (2)

- Improved land data assimilation systems :
  - EKF : SURFEX, MSC, ECMWF
  - EnKF : NILU (Met.No), US community
- Improve 2D analysis systems (SST, snow cover, screen-level variables) : OI accounting for anisotropy effects (e.g. wavelet structure functions)




# Land albedo analysis




Difference (ANA – ECO) – 13/03/2007

# Impact on forecasts




T2m forecast scores in March 2007




Differences in T2m FC+12  
13 March 2007

# Snow analysis with satellite obs



## Example from COSMO – 19/01/2006

COSMO  
OLD



IFS

COSMO  
NEW

MSG  
snow mask


Figure 2: Snow analysis on 19.01.2006. Clockwise from upper left panel: DWD product, ECMWF product, MSG derived snow mask for that day (composite map), and new product discussed in this paper. Note the large snow patch over North-Eastern Germany not present in the new product; the snow map shows bare soil over this region (in grey), but the corresponding satellite information is old due to overcast situation (not shown).

From J.M. Bettems (Meteo Swiss)


# Soil moisture analysis

SURFEX EKF (accumulated increments in root zone)

SYNOP (T2m,HU2m) –ERS (SM)



ERS (SM)




July 2006

-300 -200 -100 -50 -10 10 50 100 200 300







# Validation studies

- Complement SYNOP data (screen-level parameters and precipitation)
  - Share data from instrumented sites (over long periods of time – limited interest of data from field campaigns) – fluxes + soil variables :
    - SMOSREX, SMOSMANIA (Météo-France)
    - Lindenberg, Payerne, Cabauw
- 



# SMOSMANIA


- Soil Moisture Observing System – Meteorological Automatic Network Integrated Application
  - 12 stations at already existing AWS locations
  - Real-time soil moisture measurements –5cm –10cm –20cm –30cm
  - 1 measurement every 12 minutes



From J.C. Calvet (MF)

# Model Monitoring in COSMO

Monitoring of model near-surface and soil variables against observations  
Richard-Assmann Observatory (RAO)  
Lindenberg (52.15 N, 14.11 E)



— Observation  
— Model

